arXiv:math/0606781v1 [math.CA] 30 Jun 2006

ON A RELATION BETWEEN q-EXPONENTIAL AND θ-FUNCTION

RUIMING ZHANG

Abstract. We will use a discrete analogue of the classical Laplace method to show that for infinitely many positive integers n, the main term of the asymptotic expansion of the scaled q-exponential $(-q^{-nt+1/2}u;q)_\infty$ could be expressed in θ-function.

1. Introduction

Given an arbitrary positive real number $0<q<1$ and an arbitrary complex number a, we define[6, 10]

$$(a;q)_\infty=\prod_{k=0}^{\infty}(1-aq^k) \tag{1.1}$$

and the q-shifted factorial as

$$(a;q)_n=\frac{(a;q)_\infty}{(aq^n;q)_\infty} \tag{1.2}$$

for any integer n. Assume that $|z|<1$, the q-Binomial theorem is[6, 10]

$$\frac{(az;q)_\infty}{(z;q)_\infty}=\sum_{k=0}^{\infty}\frac{(a;q)_k}{(q;q)_k}z^k. \tag{1.3}$$

This could be seen easily from the fact that

$$\frac{(az;q)_\infty}{(z;q)_\infty}=\prod_{k=0}^{\infty}\frac{1-azq^k}{1-zq^k} \tag{1.4}$$

defines an analytic function in the region $|z|<1$. Hence it could be expanded as a power series

$$\frac{(az;q)_\infty}{(z;q)_\infty}=\sum_{k=0}^{\infty}a_kz^k \tag{1.5}$$

for $|z|<1$ with

$$a_0=1. \tag{1.6}$$

Observe that

$$\frac{(az;q)_\infty}{(z;q)_\infty}=\frac{1-az}{1-z}\frac{(azq;q)_\infty}{(zq;q)_\infty}, \tag{1.7}$$

Date: June 6, 2006.

1991 *Mathematics Subject Classification.* Primary 33D45. Secondary 33E05.

Key words and phrases. Asymptotics, θ-function, q-binomial theorem, q-exponential, discrete Laplace method.

thus we have

$$(1-z)\sum_{k=0}^{\infty} a_k z^k = (1-az)\sum_{k=0}^{\infty} a_k q^k z^k \tag{1.8}$$

or

$$\sum_{k=0}^{\infty} a_k(1-q^k)z^k = \sum_{k=1}^{\infty} a_{k-1}(1-aq^{k-1})z^k. \tag{1.9}$$

Then

$$a_k = \frac{1-aq^{k-1}}{1-q^k}a_{k-1} \tag{1.10}$$

and

$$a_k = \frac{(a;q)_k}{(q;q)_k} \tag{1.11}$$

for $k = 1, 2, \ldots$. For any complex number z, let a be a large real number with $a \gg |z|$. We replace z in (1.3) by $\frac{z}{a}$,

$$\frac{(z;q)_\infty}{(z/a;q)_\infty} = \sum_{k=0}^{\infty} \frac{(a;q)_k/a^k}{(q;q)_k} z^k. \tag{1.12}$$

Since

$$\lim_{a\to\infty} \frac{(a;q)_k}{a^k} = (-1)^k q^{k(k-1)/2}, \tag{1.13}$$

then

$$\lim_{a\to\infty} \sum_{k=0}^{\infty} \frac{(a;q)_k/a^k}{(q;q)_k} z^k = \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{(q;q)_k}(-z)^k \tag{1.14}$$

by the Lebesgue dominated convergent theorem. Similarly from (1.3)

$$\frac{1}{(z/a;q)_\infty} = \sum_{k=0}^{\infty} \frac{1}{(q;q)_k}\left(\frac{z}{a}\right)^k, \tag{1.15}$$

then

$$\begin{aligned} \lim_{a\to\infty} \frac{(z;q)_\infty}{(z/a;q)_\infty} &= \lim_{a\to\infty} \sum_{k=0}^{\infty} \frac{(z;q)_\infty}{(q;q)_k}\left(\frac{z}{a}\right)^k \\ &= (z;q)_\infty \end{aligned} \tag{1.16}$$

by the Lebesgue dominated theorem. Thus we have proved that for any complex number z

$$(z;q)_\infty = \sum_{k=0}^{\infty} \frac{q^{k(k-1)/2}}{(q;q)_k}(-z)^k. \tag{1.17}$$

Notice that

$$((1-q)z;q)_\infty = \sum_{k=0}^{\infty} \frac{(1-q)^k}{(q;q)_k} q^{k(k-1)/2}(-z)^k. \tag{1.18}$$

Since

$$\frac{1-q^k}{1-q} \geq kq^{k-1} \tag{1.19}$$

for $k \geq 1$, thus

$$\frac{(1-q)^k q^{k(k-1)/2}}{(q;q)_k} \leq \frac{1}{k!} \tag{1.20}$$

for $k \geq 1$, hence by the Lebesgue dominated theorem we have

$$\lim_{q\to 1}((1-q)z;q)_\infty = \sum_{k=0}^{\infty} \frac{(-z)^k}{k!} = e^{-z}, \tag{1.21}$$

and this is the reason why $(z;q)_\infty$ is called a q-exponential. Indeed, it is one of several q-analogues of e^z in the theory of q-series. From (1.18) and (1.20), it is also clear that

$$|((1-q)z;q)_\infty| \leq e^{|z|} \tag{1.22}$$

for any complex number z. For any nonzero complex number z, we define the theta function as

$$\theta(z;q) = \sum_{k=-\infty}^{\infty} q^{k^2/2} z^k. \tag{1.23}$$

The Jacobi's triple product formula says that[6, 10]

$$\sum_{k=-\infty}^{\infty} q^{k^2/2} z^k = (q, -q^{1/2}z, -q^{1/2}/z; q)_\infty. \tag{1.24}$$

For any positive real number t, we consider the following set

$$\mathbb{S}(t) = \{\{nt\} : n \in \mathbb{N}\}. \tag{1.25}$$

It is clear that $\mathbb{S}(t) \subset [0,1)$ and it is a finite set when t is a positive rational number. In this case, for any $\lambda \in \mathbb{S}(t)$, there are infinitely many positive integers n and m such that

$$nt = m + \lambda, \tag{1.26}$$

where

$$m = \lfloor nt \rfloor. \tag{1.27}$$

If t is a positive irrational number, then $\mathbb{S}(t)$ is a subset of $(0,1)$ with infinite elements, and it is well-known that $\mathbb{S}(t)$ is uniformly distributed in $(0,1)$. A theorem of Chebyshev[8] says that given any $\beta \in [0,1)$, there are infinitely many positive integers n and m such that

$$nt = m + \beta + \gamma_n \tag{1.28}$$

with

$$|\gamma_n| \leq \frac{3}{n}, \tag{1.29}$$

It is clear that for n large enough, we may also have

$$m = \lfloor nt \rfloor. \tag{1.30}$$

2. MAIN RESULTS

From (1.17), clearly we have

$$(-q^{-nt+1/2}u;q)_\infty = \sum_{k=0}^{\infty}\frac{q^{k^2/2-knt}}{(q;q)_k}u^k. \tag{2.1}$$

Theorem. *Given an arbitrary positive real number $0 < q < 1$ and an arbitrary nonzero complex number u, we have*

1. *For any positive rational number t, let $\lambda \in \mathbb{S}(t)$, there are infinitely many positive integers n and m such that*

$$tn = m + \lambda, \tag{2.2}$$

and for these n's and m's we have

$$(-q^{-nt+1/2}u;q)_\infty = \frac{u^m\left\{\theta(u^{-1}q^\lambda;q) + r(n)\right\}}{(q;q)_\infty q^{m^2/2+m\lambda}} \tag{2.3}$$

with

$$\begin{aligned} |r(n)| &\le \frac{3(-q;q)_\infty\theta(|u|^{-1}q^\lambda;q)}{1-q} \\ &\times \left\{q^{m/2} + \frac{q^{m^2/8}}{|u|^{\lfloor m/2\rfloor+1}}\right\}. \end{aligned} \tag{2.4}$$

for n, m and λ satisfying (2.2) with n and m are large enough.

2. *Given a positive irrational number t, for any real number $\beta \in [0,1)$, there are infinitely many positive integers n and m such that*

$$nt = m + \beta + \gamma_n \tag{2.5}$$

with

$$|\gamma_n| \le \frac{3}{n}. \tag{2.6}$$

For such n's and m's we have

$$(-q^{-nt+1/2}u;q)_\infty = \frac{u^m\left\{\theta(u^{-1}q^\beta;q) + O\left(\frac{\log n}{n}\right)\right\}}{(q;q)_\infty q^{mnt-m^2/2}}. \tag{2.7}$$

for n is large enough.

Proof. In the case that t is a positive rational number, for any $\lambda \in \mathbb{S}(t)$ and n, m are large, we have

$$\begin{aligned} (-q^{-nt+1/2}u;q)_\infty(q;q)_\infty &= \sum_{k=0}^{\infty}(q^{k+1};q)_\infty q^{k^2/2-km-k\lambda}u^k \\ &= s_1 + s_2 \end{aligned} \tag{2.8}$$

where

$$s_1 = \sum_{k=0}^{m}(q^{k+1};q)_\infty q^{k^2/2-km-k\lambda}u^k \tag{2.9}$$

and

$$s_2 = \sum_{k=m+1}^{\infty}(q^{k+1};q)_\infty q^{k^2/2-km-k\lambda}u^k. \tag{2.10}$$

In s_1 we change the summation from k to $m-k$, we have

$$\begin{aligned}
\frac{s_1 q^{m^2/2+m\lambda}}{u^m} &= \sum_{k=0}^{m}(q^{m-k+1};q)_\infty q^{k^2/2}(u^{-1}q^{\lambda})^k \\
&= \sum_{k=0}^{\infty} q^{k^2/2}(u^{-1}q^{\lambda})^k \\
&- \sum_{k=\lfloor m/2\rfloor+1}^{\infty} q^{k^2/2}(u^{-1}q^{\lambda})^k \\
&+ \sum_{k=0}^{\lfloor m/2\rfloor} q^{k^2/2}(u^{-1}q^{\lambda})^k \left((q^{m-k+1};q)_\infty - 1\right) \\
&+ \sum_{k=\lfloor m/2\rfloor+1}^{m} (q^{m-k+1};q)_\infty q^{k^2/2}(u^{-1}q^{\lambda})^k \\
&= \sum_{k=0}^{\infty} q^{k^2/2}(u^{-1}q^{\lambda})^k + s_{11} + s_{12} + s_{13}.
\end{aligned}$$

Clearly,

$$\begin{aligned}
|s_{11}+s_{13}| &\le 2\sum_{k=\lfloor m/2\rfloor+1}^{\infty} q^{k^2/2}(|u|^{-1}q^{\lambda})^k \\
&\le \frac{2q^{m^2/8}}{|u|^{\lfloor m/2\rfloor+1}}\theta(|u|^{-1}q^{\lambda};q).
\end{aligned}$$

By (1.3), for $0 \le k \le \lfloor m/2\rfloor$, we have

$$\left|(q^{m-k+1};q)_\infty - 1\right| = \frac{(-q;q)_\infty}{1-q}q^{m/2}, \tag{2.11}$$

then

$$\begin{aligned}
|s_{12}| &\le \frac{(-q;q)_\infty}{1-q}q^{m/2}\sum_{k=0}^{\infty} q^{k^2/2}(|u|^{-1}q^{\lambda})^k \\
&\le \frac{q^{m/2}(-q;q)_\infty\theta(|u|^{-1}q^{\lambda};q)}{1-q},
\end{aligned}$$

hence

$$\frac{s_1 q^{m^2/2+m\lambda}}{u^m} = \sum_{k=0}^{\infty} q^{k^2/2}(u^{-1}q^{\lambda})^k + r_1(n) \tag{2.12}$$

with

$$\begin{aligned}
|r_1(n)| &\le \frac{2(-q;q)_\infty\theta(|u|^{-1}q^{\lambda};q)}{1-q} \\
&\times \left\{q^{m/2} + \frac{q^{m^2/8}}{|u|^{\lfloor m/2\rfloor+1}}\right\}.
\end{aligned} \tag{2.13}$$

In s_2 we change the summation from k to $k+m$

$$\begin{aligned}\frac{s_2q^{m^2/2+m\lambda}}{u^m} &= \sum_{k=1}^{\infty}(q^{m+k+1};q)_\infty q^{k^2/2}(uq^{-\lambda})^k\\ &= \sum_{k=1}^{\infty} q^{k^2/2}(uq^{-\lambda})^k\\ &+ \sum_{k=1}^{\infty} q^{k^2/2}(uq^{-\lambda})^k\left[(q^{m+k+1};q)_\infty-1\right]\\ &= \sum_{k=-\infty}^{-1} q^{k^2/2}(u^{-1}q^{\lambda})^k + r_2(n).\end{aligned}$$

By the binomial theorem, for $k \ge 1$

$$\left|(q^{m+k+1};q)_\infty - 1\right| \le \frac{q^{m+2}(-q^3;q)_\infty}{1-q}, \tag{2.14}$$

then

$$|r_2(n)| \le \frac{q^{m+2}(-q^3;q)_\infty\theta(|u|^{-1}q^{\lambda};q)}{1-q}. \tag{2.15}$$

Thus we have proved that

$$(-q^{-nt+1/2}u;q)_\infty = \frac{u^m\left\{\theta(u^{-1}q^{\lambda};q)+r(n)\right\}}{(q;q)_\infty q^{m^2/2+m\lambda}} \tag{2.16}$$

with

$$\begin{aligned}|r(n)| &\le \frac{3(-q;q)_\infty\theta(|u|^{-1}q^{\lambda};q)}{1-q}\\ &\times \left\{q^{m/2}+\frac{q^{m^2/8}}{|u|^{\lfloor m/2\rfloor+1}}\right\}.\end{aligned} \tag{2.17}$$

for n, m and λ satisfying (2.2) with n and m are large enough.

In the case that t is a positive irrational number, for any real number $\beta \in [0,1)$, when n and m are large enough satisfying (2.5) and (2.6), we have

$$\beta + \gamma_n \ge -1, \tag{2.18}$$

for these integers n, we take

$$\nu_n = \left\lfloor -\frac{\log n}{\log q}\right\rfloor, \tag{2.19}$$

then we have

$$\begin{aligned}(-q^{-nt+1/2}u;q)_\infty(q;q)_\infty &= \sum_{k=0}^{\infty}(q^{k+1};q)_\infty q^{k^2/2-km-k\beta-k\gamma_n}u^k\\ &= s_1+s_2,\end{aligned} \tag{2.20}$$

with

$$s_1 = \sum_{k=0}^{m}(q^{k+1};q)_\infty q^{k^2/2-km-k\beta-k\gamma_n}u^k \tag{2.21}$$

and

$$s_2 = \sum_{k=m+1}^{\infty} (q^{k+1};q)_\infty q^{k^2/2-km-k\beta-k\gamma_n} u^k. \tag{2.22}$$

In s_1, we change summation from k to $m-k$,

$$\begin{aligned}
\frac{s_1 q^{mnt-m^2/2}}{u^m} &= \sum_{k=0}^{m} (q^{m-k+1};q)_\infty q^{k^2/2}(u^{-1}q^{\beta+\gamma_n})^k \\
&= \sum_{k=0}^{\infty} q^{k^2/2}(u^{-1}q^{\beta})^k \\
&- \sum_{k=\nu_n+1}^{\infty} q^{k^2/2}(u^{-1}q^{\beta})^k \\
&+ \sum_{k=0}^{\nu_n} q^{k^2/2}(u^{-1}q^{\beta})^k \left(q^{k\gamma_n}-1\right) \\
&+ \sum_{k=0}^{\nu_n} q^{k^2/2}(u^{-1}q^{\beta+\gamma_n})^k \left\{(q^{m-k+1};q)_\infty - 1\right\} \\
&+ \sum_{k=\nu_n+1}^{m} q^{k^2/2}(u^{-1}q^{\beta+\gamma_n})^k (q^{m-k+1};q)_\infty \\
&= \sum_{k=0}^{\infty} q^{k^2/2}(u^{-1}q^{\beta})^k + s_{11} + s_{12} + s_{13} + s_{14},
\end{aligned}$$

thus there exists some constant $c_{11}(q,u)$ such that

$$\begin{aligned}
|s_{11}+s_{14}| &\le 2\sum_{k=\nu_n+1}^{\infty} q^{k^2/2}\,|u|^{-k} \\
&\le c_{11}(q,u)\frac{\log n}{n}
\end{aligned}$$

for n large enough, and there exists some constant $c_{12}(q,u)$ such that

$$|s_{12}+s_{13}| \le c_{12}(q,u)\frac{\log n}{n},$$

for n large enough. Thus for n, m and β satisfying (2.5) and (2.6) and n is large enough, there exists some constant $c_1(q,u)$ such that

$$\frac{s_1 q^{mnt-m^2/2}}{u^m} = \sum_{k=0}^{\infty} q^{k^2/2}(u^{-1}q^{\beta})^k + e_1(n) \tag{2.23}$$

with

$$|e_1(n)| \le c_1(q,u)\frac{\log n}{n}. \tag{2.24}$$

Similarly, in s_2 we change summation from k to $k+m$, and we could show that there for n, m and β satisfying (2.5) and (2.6) and n is large enough, there exists some constant $c_2(q,u)$ such that

$$\frac{s_2 q^{mnt-m^2/2}}{u^m} = \sum_{k=-\infty}^{-1} q^{k^2/2}(u^{-1}q^{\beta})^k + e_2(n) \tag{2.25}$$

with

$$|e_2(n)| \le c_2(q,u)\frac{\log n}{n}. \tag{2.26}$$

Thus we have

$$(-q^{-nt+1/2}u;q)_\infty = \frac{u^m\left\{\theta(u^{-1}q^{\beta};q) + O\left(\frac{\log n}{n}\right)\right\}}{(q;q)_\infty q^{mnt-m^2/2}} \tag{2.27}$$

for n, m and β satisfying (2.5) and (2.6) and n is large enough. ☐

Conclusion. The phenomenon demonstrated here is universal for class of entire basic hypergeometric functions, the general proof for this phenomenon will be published elsewhere. In a separated work, we have proved the same phenomenon happens with the Ramanujan's entire function $A_q(z)$, or the q-Airy function.

REFERENCES

[1] N. I. Akhiezer, *The Classical Moment Problem and Some Related Questions in Analysis*, English translation, Oliver and Boyed, Edinburgh, 1965.

[2] N. M. Atakishiyev, *A simple difference realization of the Heisenberg qalgebra*, J. Math. Phys. 35 (1994), 3253–3260.

[3] J. S. Christiansen, *The moment problem associated with the Stieltjes–Wigert polynomials*, J. Math. Anal. Appl. 277 (2003), 218–245.

[4] P. Deift, *Orthogonal Polynomials and Random Matrices: a Riemann-Hilbert Approach*, American Mathematical Society, Providence, 2000.

[5] P. Deift, T. Kriecherbauer, K. T-R. McLaughlin, S. Venakides, and X. Zhou, *Strong asymptotics of orthogonal polynomials with respect to exponential weights*, Comm. Pure Appl. Math., 52 (1999), 1491–1552.

[6] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, second edition Cambridge University Press, Cambridge, 2004.

[7] W. K. Hayman, *On the zeros of a q-Bessel function*, Contemporary Mathematics, volume 382, American Mathematical Society, Providence, 2005, 205–216.

[8] Hua Loo Keng, *Introduction to Number Theory*, Springer-Verlag, Berlin Heidelberg New York, 1982.

[9] M. E. H. Ismail, *Asymptotics of q-orthogonal polynomials and a q-Airy function*, IMRN 2005 No 18 (2005), 1063–1088.

[10] M. E. H. Ismail, *Classical and Quantum Orthogonal Polynomials in one Variable*, Cambridge University Press, Cambridge, 2005.

[11] M. E. H. Ismail and X. Li, *Bounds for extreme zeros of orthogonal polynomials*, Proc. Amer. Math. Soc. 115 (1992), 131–140.

[12] M. E. H. Ismail and D. R. Masson, *q-Hermite polynomials, biorthogonal rational functions*, Trans. Amer. Math. Soc. 346, (1994), 63–116.

[13] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta, Y. Yamada, *Hypergeometric solutions to the q-Painlevé equations*, Internat. Math. Res. Notices 47 (2004), 2497–2521.

[14] R. Koekoek and R. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogues*, Reports of the Faculty of Technical Mathematics and Informatics no. 98-17, Delft University of Technology, Delft, 1998.

[15] W.-Y. Qiu and R.Wong, *Uniform asymptotic formula for orthogonal polynomials with exponential weight*, SIAM J. Math. Anal., 31, (2000), 992–1029.

[16] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers* (Introduction by G. E. Andrews), Narosa, New Delhi, 1988.

[17] G. Szego, *Orthogonal Polynomials*, Fourth Edition, Amer. Math. Soc., Providence, 1975.

[18] Z. Wang and R. Wong, *Uniform asymptotics for the Stieltjes-Wigert polynomials: the Riemann-Hilbert approach*, to appear.
[19] E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, fourth edition, Cambridge University Press, Cambridge, 1927.

School of Mathematics, Guangxi Normal University, Guilin City, Guangxi 541004, P. R. China.
E-mail address: `ruimingzhang@yahoo.com`